



ISSN: 2456-4419

Impact Factor: (RJIF): 5.18

Yoga 2018; 3(1): 545-547

© 2018 Yoga

www.theyogicjournal.com

Received: 17-11-2017

Accepted: 18-12-2017

**Dr. Parwinder Singh**

Assistant Professor, Department  
of Sports, RIMT University  
Mandi Gobindgarh, Punjab,  
India

**Dr. Ashok Kumar**

Associate Professor, Hod  
Department of Sports Science,  
Punjabi University Patiala,  
Punjab, India

**Harinder Singh**

Assistant Director of Physical  
Education, RIMT University  
Mandi Gobindgarh, Punjab,  
India

## A study on body weight and body composition of junior free style wrestlers

**Dr. Parwinder Singh, Dr. Ashok Kumar and Harinder Singh Chatha**

### Abstract

**Aim:** The aim of this study was to examine weight and the body composition of different weight categories of junior free style male wrestlers.

**Method:** One hundred fifty (N=150) male junior free style wrestlers were participated as subjects and they were further divided into five groups according to their weight categories, each group was comprised of thirty wrestlers. Body fat percentage was estimated by using Durnin and Womersley equation and Body weight was measured with weighing machine.

**Results:** A statistical significant positive correlation was found between weight and BMI, skinfold thickness and %BF.

**Conclusion:** It is concluded that the free style wrestling is a weight classified sport, therefore the correlation of weight of the wrestlers with body composition is important. The results of the present study also demonstrated the effect of age, height, BMI, and BF % on the weight of junior free style wrestlers.

**Keywords:** Free style, wrestling, skinfold, anthropometry, strength, fat percentage

### Introduction

A sport of wrestling has a strong tradition that precedes the first Olympic festival in 776 B. C., when Zeus wrestled Kronas for the possession of the earth (Gallagher, 1951). Since then the sport of wrestling continues to grow in popularity because of the discipline and mental toughness it requires to be successful in the sport. Unfortunately, the sport has also been associated with the stigma of "cutting weight" and the practices that accompany the process of competing at designated weight classes. Like the sports of judo, boxing, and competitive weight lifting, wrestling requires its athletes to compete at specific weights or weight classifications. Typically, these weight classifications differ by approximately 7-11 pounds depending on age and style of wrestling. It is common knowledge in present day of wrestling that wrestlers compete in weight classes below their "normal" weight. The purpose of this practice is to gain advantages in strength, speed, and leverage over their opponents (Steen & Brownell, 1990). The changes in regulation of wrestling have forced several modifications in the fitness requirements of successful wrestlers, which as a result caused an evolution in the training methods (Yoon 2002; Horswill 1992; Sharratt et. al., 1986). Wrestling has been described as an intermittent physical event which produces great strength and muscle power demands of both the upper and lower body (Hubner-Wozniak et. al. 2004; Kraemer et. al. 2001; Horswill et. al. 1992 & 1989; Sharratt et. al. 1986). They generally want to minimize the body fat level and the total body weight without losing their body strength and power (Yoon, 2002). However, no relation was shown between the percentage of fat mass (%FM) and the level of wrestling success (Yoon, 2002; Horswill 1992). The aim of the present study was to observe the body composition and body weight in different weight categories of junior free style wrestlers. The present investigation was to study the relationship of body composition and body weight in different weight categories of junior free style wrestlers.

### Material and Methods

The study was conducted on 150 male junior free style wrestlers (age between 18-20 years) of five different groups i.e. Group 1- 46-50 kg junior free style wrestlers, Group 2- 55kg junior

### Correspondence

**Dr. Parwinder Singh**

Assistant Professor, Department  
of Sports, RIMT University  
Mandi Gobindgarh, Punjab,  
India

free style wrestlers, Group 3- 60kg junior free style wrestlers, Group 4 – 66kg junior free style wrestlers, Group 5 – 74kg junior free style wrestlers. The aim of the study was explained to each participant and signed informed consent was obtained from the participants. The body composition of junior free style wrestlers was estimated as per the method described by Durmin and Womersley (1974). BMI can be categorized according to the WHO (Laquatra, 2004). Body weight was measured with weighing machine. Karl Pearson’s coefficient

of correlation was used to find the relationship among age, weight, BMI and body composition variables of free style junior wrestler in five groups according to their weight. To determine the differences between the mean of the various variables among different groups, one way Analysis of Variance (ANOVA) was used. Scheffe Post Hoc test was also used to identify the location of significant differences among the different groups. The level of significance was  $p < 0.05$ .

**Table 1:** Mean± SD of age, height, weight, BMI and % body fat of different groups

Variable(s)	Group 1 (n=30)	Group 2 (n=30)	Group 3 (n=30)	Group 4 (n=30)	Group 5 (n=30)	Total (n=150)
Age (year)	18.13±0.77	18.73±0.69	18.53±0.68	18.50±0.68	18.67±0.47	18.51±0.65
Height (cm)	166.69±2.62	168.00±2.62	168.46±3.24	168.95±2.79	171.80±4.37	168.78±3.12
Weight (kg)	51.50±1.33	55.70±1.41	61.50±0.88	66.33±1.40	74.25±3.22	61.85±1.64
BMI	18.54±0.65	19.74±.77	21.69±0.90	23.25±0.85	25.18±1.43	21.68±0.92
% Body fat	9.10±1.67	10.32±2.13	11.92±1.43	12.71±2.28	14.88±3.06	11.78±2.11

**Results**

Table 1 shows that the mean age of group1, group2, group3, group4 and group5 was 18.13±.77 year, 18.73±.69 year, 18.53±.68 year, 18.50±.68 year and 18.67±.47 year. The mean height of the subjects of group1, group2, group3, group4 and group5 was 166.69±2.62cm year, 168.00±2.62cm, 168.46±3.24cm, 168.95±2.79cm and 171.80±4.37cm. The mean weight of the subjects of group1, group2, group3, group4 and group5 was 51.50±1.33kg, 55.70±1.41kg, 61.50±.88 kg, 66.33±1.40 kg and 74.25±3.22 kg. The mean

body mass index of the subjects of group1, group2, group3, group4 and group5 was 18.54±.65 Kg/m<sup>2</sup>, 19.74±.77Kg/m<sup>2</sup>, 21.69±.90Kg/m<sup>2</sup>, 23.25±.85Kg/m<sup>2</sup> and 25.18±1.43Kg/m<sup>2</sup> respectively. The mean percent fat of the subjects of group1, group2, group3, group4 and group5 was 9.10±1.67%, 10.32±2.13%, 11.92±1.43%, 12.71±2.28% and 14.88±3.06 % . Further, the analysis of variance (Table 2) revealed that the variance in the mean values of age, height, weight, BMI and % body fat of junior free style wrestlers among different groups was statistical significant.

**Table 2:** Analysis of variance of age, height, weight, BMI and % body fat among different groups

Variable(s)	Groups	Sum of Squares	Mean Square	F	Sig.
Age	Between Groups	6.50	1.627	3.63	.008
	Within Groups	64.96	.448		
Height	Between Groups	427.42	106.855	10.43	.000
	Within Groups	1484.95	10.241		
Weight	Between Groups	9562.39	2390.600	706.97	.000
	Within Groups	490.31	3.381		
Body mass index	Between Groups	849.94	212.485	230.09	.000
	Within Groups	133.90	.923		
Percentage Fat	Between Groups	594.57	148.644	30.86	.000
	Within Groups	698.32	4.816		

\*significant at the 0.05 level

Table 3 shows a positive statistical significant relationship among percent body fat, height, and weight of combined (N=150) groups of junior free style wrestlers. A positive

statistical significant relationship was also found between weight, percent body fat and BMI (Table 3).

**Table 3:** Correlation among age, height, weight, BMI and % body fat variables of combined group (N=150)

Variable(s)	Height	weight	Body mass index	Percentage fat
Age	.076	.168*	.160	.080
Height		.483**	.167*	.257**
Weight			.944**	.722**
BMI				.717**

\*significant at the 0.05 level

**Discussion**

The results of the present study shows that the body fat percentage of the wrestlers of different groups were in the acceptable range according to the WHO (Laquatra, 2004). In other words, we can say no obesity was observed in different group of junior free style wrestlers. The maximum body fat percentage was observed in group 5 (higher weight category) and minimum in group1 (lower weight) wrestlers. Thus, a trend of increased in body fat percentage from lower body weight group1 (46 kg- 50kg) to higher body weight group5

(74kg) of wrestlers was observed. The maximum lean body mass percentage was observed in group1 and minimum lean body mass percentage was observed in group 5. Thus, a trend of decrease in lean body mass percentage was observed from lower body weight group1 (46-50kg) to higher body weight group5 (74kg). Saygin (2014) also reported that the percentage body fat has different values in all three classes like lightweight, middle weight and heavy weight wrestlers. According to Saygin (2014) the maximum body fat was observed in heavyweight wrestlers when compared to

lightweight wrestlers and middleweight wrestlers. Franchini *et al.* (2014) also reported that the highest skinfold thickness values were observed in heavyweight judo athletes as compared to lightweight and middleweight judo athletes, in other words, heavy weight judo athletes has more fat percentage than lightweight and middle weight. The mean value of various anthropometric variables body mass index was increased from lower body weight (group1) to higher body weight (group 5). Selda (2014) also reported that there was a significant difference in the body mass index of different weight categories of greco roman and free style wrestlers and the body mass index was higher in heavy weight greco roman and free style wrestlers as compared to the lower and middle weight greco roman and free style wrestlers. Selda (2014) reported that both heavy weight greco roman and free style group wrestlers was taller than the lower weight and middle weight wrestlers and a similar trend of body height was also found in the present study (i.e. junior free style wrestlers).

### Conclusion

From the results of the present study, it is concluded that the body fat percentage of the wrestlers of different groups was in the acceptable range i.e. obesity was not observed in them. Further, it is concluded that the wrestlers of lower body weight category (group1) were better in anthropometric variables than higher body weight category (group 5) like body fat percentage, body mass index. A positive relationship was also observed between weight, age, height, BMI and % body fat,.

### References

1. Durnin JV, Womersley J. Body fat assessed from total body density and its estimation from skinfold thickness: measurements on 481 males and females aged from 16 to 72 years. *Br J Nutr.* 1974; 32:77-97.
2. Franchini Emerson, Katarzyna Sterkowicz-Przybycien and Monica Yuri Takito. Anthropometrical Profile of Judo Athletes: Comparative Analysis between Weight Categories. *Int. J. Morphol.* 2014; 32(1):36-42.
3. Gallagher EC. *Wrestling (Revised Edition ed.)*. New York: The Ronald Press Company. 1951.
4. Horswill CA. Applied physiology of amateur wrestling. *Sports Med* 1992; 14:114-143.
5. Horswill CA, Miller JE, Scott JR, Smith CM, Welk G, Von Handel P, Anaerobic and aerobic power in arms and legs of elite senior wrestlers *Int J Sports Med.* 1992; 13, 558-561.
6. Horswill CA, Scott JR, Galea P. Comparison of maximum aerobic power, maximum anaerobic power, and skinfold thickness of elite and nonelite junior wrestlers. *Int J Sports Med.* 1989; 10:165-168.
7. Hubner Wozniak E, Kosmol A, Lutoslawska G, Bem EZ. Anaerobic performance of arms and legs in male and female free style wrestlers. *J Sci Med Sport.* 2004; 7:473-480.
8. Kraemer WJ, Fry AC, Rubin MR, Triplett-mcbride T, Gordon SE, Koziris LP *et al.* Physiological and performance responses to tournament wrestling. *Med Sci Sports Exerc.* 2001; 33:1367-1378.
9. Laquatra I. Nutrition for weight management. In Krause s Food, Nutrition, and Diet Therapy. Ed. by Mahan, LK. and Escott-Stumps, S. 11th ed. Saunders Company, Philadelphia, Pennsylvanian. 2004; 558-593.
10. Saygin O. Examination of Some Physical, Hematological

Parameters and Iron Status of Greco-Roman Wrestlers in the Age Category of Cadets by Weight Classes *Anthropologist.* 2014; 18(2):25-334.

11. Selda Basara, IremDuzguna, Nevin Atalay Guzela, Ibrahim Cicioglu and Bulent Celikc () Differences in strength, flexibility and stability in freestyle and Greco-Roman wrestlers. *Musculoskeletal Rehabilitation.* 2014; 27 321-330.
12. Sharratt MT, Taylor AW, Song TM. A physiological profile of elite Canadian freestyle wrestlers. *Can J Appl Sport Sci.* 1986; 11:100-105.
13. Steen SN, Brownell KD. Patterns of weight loss and regain in wrestlers: has the tradition changed. *Med Sci Sports Exercise.* 1990; 22(6):762-768.
14. Yoon J. Physiological profiles of elite senior wrestlers. *Sports Med.* 2002; 32:225-233.