



ISSN: 2456-4419

Impact Factor: (RJIF): 5.18

Yoga 2020; 5(2): 14-15

© 2020 Yoga

www.theyogicjournal.com

Received: 15-05-2020

Accepted: 19-06-2020

Nagaraj GP

Physical Education Teacher,
Diploma Agriculture College
Katalagere Channagiri Taluk,
Davanagere, Karnataka, India

Dr. SK Manjunatha

Assistant Professor, Department
of Physical Education, College of
Horticultural, Mudigere,
Karnataka, India

Eshwara KA

Assistant Professor, Department
of Physical Education, College of
Forestry, Ponnampet,
Karnataka, India

Corresponding Author:

Nagaraj GP

Physical Education Teacher,
Diploma Agriculture College
Katalagere Channagiri Taluk,
Davanagere, Karnataka, India

Effects of physical activity on chronic kidney diseases

Nagaraj GP, Dr. SK Manjunatha and Eshwara KA

Abstract

Chronic kidney disease also known as chronic renal disease. Chronic kidney diseases of long duration and generally slow progression it is such as heart diseases stroke, cancer, chronic respiratory diseases and diabetes. The symptoms of kidney disease generally unwell feelings reduce aptitude and more sweating on the night bed. Those who has high blood pressure diabetes and blood relative problem, more chance to be chronic disease. Physical activity is very necessary for patient of the chronic kidney diseases. Yoga and therapy is very beneficial for the chronic kidney and heart diseases also. Types of Kidney Diseases they are 1. Diabetes, 2. High blood pressure (hyper tension), 3. Hereditary, 4. Kidney stone, 5. Drug and toxins, 6. Cancers 7. HIV Infections and 8. Smoking. 1st Stage Slightly diminished function GFR (>90 mL/min/1.73 m²). 2nd Stage Mild reduction in GFR (60-89 mL/min/1.73 m²). 3rd Stage Moderate reduction in GFR (30-59 mL/min/1.73 m²). 4th Stage Severe reduction in GFR (15-29 mL/min/1.73 m²). 5th Stage Established kidney failure (GFR <15 mL/min/1.73 m²). Physical activity is plays important role to prevent chronic kidney diseases. If we can do per day physical activity, therapy and yoga, avoid from this diseases. Mostly related to the kidney and heart risk for developing the coronary artery diseases because of cholesterol level, diabetes and stress.

Keywords: Physical activity, chronic kidney diseases

Introduction

Chronic kidney disease also known as chronic renal disease. Chronic kidney diseases of long duration and generally slow progression it is such as heart diseases stroke, cancer, chronic respiratory diseases and diabetes. The symptoms of kidney disease generally unwell feelings reduce aptitude and more sweating on the night bed. Those who has high blood pressure diabetes and blood relative problem, more chance to be chronic disease. Physical activity is very necessary for patient of the chronic kidney diseases. Yoga and therapy is very beneficial for the chronic kidney and heart diseases also.

Function of kidney

1. Remove waste products.
2. Regulate the chemical level in the blood
3. Kidney produced some hormones like
 - a. Kidney stimulates red blood cells production.
 - b. Regulate blood pressure
 - c. Control calcium metabolism

Chronic Diseases

Mostly related to the kidney and heart risk for developing the coronary artery diseases because of cholesterol level, diabetes and stress.

Kidney Diseases

It is disorders that affected kidney”

Types of Kidney Diseases

1. Diabetes
2. High blood pressure (hyper tension)
3. Hereditary

4. Kidney stone
5. Drug and toxins
6. Cancers
7. HIV infections
8. Smoking

Symptoms

1. High blood pressure
2. Fatigue and weakness
3. loss appetite
4. Nausea and vomiting
5. Swelling in lungs
6. Puffiness around the eyes
7. Headaches
8. Numb ness in the feet or hand
9. Disturbed sleep
10. Rest less legs syndrome
11. Chest pain
12. Bone pain
13. Poor blood clotting

Stages of the patients

1. **1st Stage:** - Slightly diminished function GFR (>90 mL/min/1.73 m²).
2. **2nd Stage:** - Mild reduction in GFR (60-89 mL/min/1.73 m²).
3. **3rd Stage:** - Moderate reduction in GFR (30-59 mL/min/1.73 m²).
4. **4th Stage:** - Severe reduction in GFR (15-29 mL/min/1.73 m²).
5. **5th Stage:** - Established kidney failure (GFR <15 mL/min/1.73 m²).

Glomerular Filtration Rate (Gfr)

As kidney disease progresses, GFR falls.

1. The normal GFR in men 100-140 ml/min.
2. In women GFR 85-115 ml/min.
3. GFR may be calculated from the amount of waste products in the 24-hour. It decreased in most people with age.

Prevention

- a. Keep your B.P. bellow 130/80.
- b. Maintain your Blood sugar level
- c. Maintain your cholesterol level
- d. Control smoking
- e. In take low fatty diet
- f. Changes in life style
- g. Avoid medications

Treatment

Self-care at home

1. Protein restriction
2. Salt restriction
3. Potassium restriction
4. Phosphors restriction
5. Fluid in take
6. Stop smoking
7. Lose excess weight

Medical treatment

1. Urine test
2. Blood test
3. Electrolyte level in acid base balance
4. Estimated GFR

5. Blood cell counts
6. Biopsy

Prevention through physical activity

Physical activity is plays important role to prevent chronic kidney diseases. If we can do per day physical activity, therapy and yoga, avoid from this diseases.

New Data

1. In India 13 people out of hundred has chronic diseases.
2. 40% of kidney failure people have diabetes in the world.
3. 50% kidney failure people with diabetes in India.
4. In Chennai 3 to 4% young (20-25 years) have hypertension.
5. 50% people (50-60 years) have high B.P. in Chennai.
6. 48% people (50-60 years) have high B.P. in Delhi.
7. 7000 people have Vic timed with chronic diseases in India

Conclusion

Physical activity is plays important role to prevent chronic kidney diseases. If we can do per day physical activity, therapy and yoga, avoid from this diseases. Mostly related to the kidney and heart risk for developing the coronary artery diseases because of cholesterol level, diabetes and stress.

References

1. Association between recombinant human erythropoietin and quality of life and exercise capacity of patients receiving haemodialysis. Canadian Erythropoietin Study Group. *BMJ*. 1990; 300(6724):573-578. [PMC free article] [PubMed]
2. European best practice guidelines for the management of anemia in patients with chronic renal failure. *Nephrology Dialysis Transplantation*. 1999; 14(15):1-50. [PubMed]
3. European Best Practice Guidelines for Haemodialysis. *Nephrology Dialysis Transplantation*. 2002; 17(7):1-111. [PubMed]
4. European best practice guidelines for the management of anemia in patients with chronic renal failure. *Nephrology Dialysis Transplantation*. 2004; 19(2):1. [PubMed]
5. Higgins J, Green S, editors. *Cochrane Handbook for Systematic Reviews of Interventions 5.0.2*. The Cochrane Collaboration; 2009. [updated September 2009] Available from: www.cochrane-handbook.org.
6. Abu-Alfa AK, Cruz D, Perazella MA, Mahnensmith RL, Simon D, Bia MJ. ACE inhibitors do not induce recombinant human erythropoietin resistance in hemodialysis patients. *American Journal of Kidney Diseases*. 2000; 35(6):1076-1082. [PubMed]
7. Adamson JW. Hyporesponsiveness to erythropoiesis stimulating agents in chronic kidney disease: the many faces of inflammation. *Advances in Chronic Kidney Disease*. 2009; 16(2):76-82. [PubMed]
8. Adhikary L, Acharya S. Efficacy of IV iron compared to oral iron for increment of haemoglobin level in anemic chronic kidney disease patients on erythropoietin therapy. *JNMA; Journal of the Nepal Medical Association*. 2011; 51(183):133-136. [PubMed]
9. Agarwal R, Davis JL, Hamburger RJ. A trial of two iron-dextran infusion regimens in chronic hemodialysis patients. *Clinical Nephrology*. 2000; 54(2):105-111. [PubMed]